

ASTHMA SCHOOL MEDICATION PLAN



Student Name: _____

Birthdate: _____

School Name: _____

Grade/Rm. _____

Emergency Contact Information and Parent / Guardian Information:

Parent / Guardian-1 (name / relationship): _____

Phone (H) _____ Tel (W) _____


Parent / Guardian-2 (name / relationship): _____

Phone (H) _____ Tel (W) _____

Healthcare Provider _____ Phone: _____

Asthma Specialist: _____ Phone: _____

Emergency contact if other than above (name/relationship): _____ Phone: _____



Student
Photo

Diagnosis / Reason for Medication: Asthma Triggers to Avoid for student while at school:	<input type="checkbox"/> Asthma	<input type="checkbox"/> Other:
	<input type="checkbox"/> Smoke / fumes	<input type="checkbox"/> Animal
	<input type="checkbox"/> Mold Spores	<input type="checkbox"/> Dust Mite
	Other: _____	

YES / NO: Student is required to have quick relief asthma medication at school to provide rapid relief of asthma symptoms if needed: cough, chest tightness, wheezing, trouble breathing, shortness of breath

YES / NO: Student is required to use quick relief asthma medication BEFORE gym or other exercise to prevent exercise induced bronchospasm from asthma

YES / NO: Student is required to take daily asthma control medication at school as directed

Medication Information	
Name of Medication	YES / NO: Albuterol _____ Other: _____
Form of Medication	<input type="checkbox"/> Inhaler <input type="checkbox"/> Nebulizer <input type="checkbox"/> Dry Powder Inhaler <input type="checkbox"/> Liquid <input type="checkbox"/> Pill / Capsule
Dosage of Medication	Number of puffs _____ Other: _____
Other instructions:	YES / NO: Inhaler MUST be used with a spacer (valved holding chamber) for administration YES / NO: Please maintain a written record (Log) of all doses: YES / NO
When to administer dose	YES / NO: 5-15 minutes before gym, recess, or exercise to prevent exercise induced bronchospasm YES / NO: As needed for FAST RELIEF of chest tightness, shortness of breath, wheezing or prolonged cough or other asthma symptoms. A total of 3 doses can be given within an 8 hour interval YES / NO: Daily at _____ AM / PM for daily asthma control (long term prevention)
Repeat Dose	_____: DO NOT REPEAT the dose _____: Repeat dose one time if symptoms not gone 10 minutes after first dose AND repeat dose every 3-4 hours IF symptoms RECUR during the school day
When to call Child's Parent	If after 2 consecutive doses (2-4 puffs per dose) are given and there is no improvement in symptoms, please seek further medical attention and call parent
When to call Child's Physician	If you have concerns or questions about the student's medication or disease
Asthma Emergency	The following are possible signs of an asthma emergency: •Difficulty breathing, walking, or talking •Blue or gray discoloration of the lips or fingernails •Failure of medication to reduce worsening symptoms.
The steps that should be taken: •Activate the emergency medical system in your area. Call 911. •Call Parent/Guardian and/or Healthcare Provider	

Supervision of Medication	<input type="checkbox"/> Student is permitted to carry medication and self-administer with no supervision		
	<input type="checkbox"/> Student MAY self-administer medication BUT supervision is required for all doses		
	<input type="checkbox"/> Student requires trained assistance to administer all doses		
Expected Normal side effects:	<input type="checkbox"/> None	<input type="checkbox"/> Fast heartbeat, tremor, hyper-activity	Other: _____
	<input type="checkbox"/> None	<input type="checkbox"/> Refrigerate	Other _____
Storage Requirements			Other: _____
START Date to begin Medication	When school receives form	Other: _____	
STOP Date to discontinue Medication	End of school year	Other: _____	
Instructions for proper use of medicine are attached to this form		<input type="checkbox"/> YES	<input type="checkbox"/> NO

PLEASE COMPLETE SECTION BELOW FOR STUDENT PERMISSION TO CARRY INHALER

*******SELF-MEDICATION FOR ASTHMA INHALERS*****
Authorization (In accordance with ORC 3313.716/3313.14)**

Who keeps the bronchodilator inhaler at school?

School policy restricting possession of medication by students is insufficient grounds for preventing students with sufficient maturity from retaining possession of their bronchodilator inhaler. Such policies, when enforced, delay appropriate treatment and restrict activities unnecessarily. The decision regarding sufficient maturity of the student to be responsible for appropriate inhaler use is an individual one to be made by the parents in consultation with their physician. The inhalers pose no abuse potential or other danger to classmates. While restrictions on bronchodilator inhaler possession may be necessary for the youngest students, it constitutes unreasonable interference with the student's medical care for school personnel to unilaterally restrict possession of bronchodilator inhalers by students judged by parents and physician to have sufficient maturity to use the device appropriately. Possession of the bronchodilator inhaler by the student also promotes earlier use that decreases the risk of requiring emergency care from rapidly progressive asthma, which on rare occasion can cause hypoxia, brain damage, and death. Discussion among parents, physician, and school personnel can determine whether school-supervised administration would improve or deter compliance.

Please check if STUDENT is permitted by healthcare provider to CARRY an inhaler and SELF-MEDICATE at school.

Student Agreement and Signature:

I, _____, agree that I will:

Never allow another student to use my medication.

Keep my medication with me at all times.

Go to the School Clinic, accompanied by someone, when I used my rescue inhaler and continue to have symptoms.

Follow school policy and my medical provider's instructions as outlined in my Asthma Medication Plan.

Student Signature: _____ Date: _____

Signature of Parent/Guardian _____ Date _____

Signature of Prescriber _____ Date _____

Copies must be provided to the principal and to the nurse.

